INDEX TO AUTHORS

AARONSON, STUART A. See STEPHENSON and AARONSON, 503

ALEXANDER, P. See GRANT, CURRIE, and ALEXANDER, 150

ALKAN, SEFIK S., WILLIAMS, E. BRADY, NITECKI, DANUTE E., and GOODMAN, JOEL W. Antigen recognition and the immune response. Humoral and cellular immune responses to small mono- and bifunctional antigen molecules, 1228

ALPER, CHESTER A., BOENISCH, THOMAS, and WATSON, LILLIAN. Genetic polymorphism in human glycine-rich beta-glycoprotein, 68

ALVARES, ALVITO P., LEIGH, SUSAN, COHN, JONATHAN, and KAPPAS, ATTALLAH. Lead and methyl mercury. Effects of acute exposure on cytochrome P-450 and the mixed function oxidase system in the liver, 1406

AMOS, BERNARD. See BERKE, SULLIVAN, and AMOS, 1334

ANDERSON, R. E., SPRENT, J., and MILLER, J. F. A. P. Cell-to-cell interaction in the immune response. VIII. Radiosensitivity of thymus-derived lymphocytes, 711

ANDERSSON, BIRGER. Studies on antibody affinity at the cellular level. Correlation between binding properties of secreted antibody and cellular receptor for antigen on immunological memory cells, 312

ANDRÉ-SCHWARTZ, JANINE. See GLEICHMANN, GLEICHMANN, ANDRÉ-SCHWARTZ, and SCHWARTZ, 516

AOKI, TADAO, and TAKAHASHI, TOSHITADA. Viral and cellular surface antigens of murine leukemias and myelomas. Serological analysis by immunoelectron microscopy, 443

ARROYAVE, CARLOS M. See KOLB, HAXBY, ARROYAVE, and MÜLLER-EBERHARD, 549

ASOFSKY, RICHARD. See CANTOR and ASOFSKY, 764

—. See LAWTON, ASOFSKY, HYLTON, and COOPER, 277

—. See PIERCE, SOLLIDAY, and ASOFSKY, 675, 698

—. See TIGELAAR and ASOFSKY, 1059

ATKINS, ELISHA, FELDMAN, JOSEPH D., FRANCIS, LORRAINE, and HURSH, ELLEN. Studies on the mechanism of fever accompanying delayed hypersensitivity. The role of the sensitized lymphocyte, 1113

ATWELL, JOHN L. See MARCHALONIS, CONE, and ATWELL, 956

AUSTEN, K. FRANK. See KAPLAN, KAY, and AUSTEN, 81

AXELRAD, M. A. See HASKILL, ELLIOTT, KERBEL, AXELRAD, and EIDINGER, 1410

BACH, F. H. See WIDMER and BACH, 1204

BASTEN, A., MILLER, J. F. A. P., SPRENT, J., and PYE, J. A receptor for antibody on B lymphocytes. I. Method of detection and functional significance, 610

—, WARNER, N. L., and MANDEL, T. II. Immunochemical and electron microscopy characteristics, 627

BAUR, SONIA. See SHERR, BAUR, GRUNDKE, ZELIGS, ZELIGS, and UHR, 1392

BECKER, ELMER L. The relationship of the chemotactic behavior of the complement-derived factors, C3a, C5a, and C567, and a bacterial chemotactic factor to their ability to activate the proesterase 1 of rabbit polymorphonuclear leukocytes, 376

BEER, ALAN E., BILLINGHAM, R. E., and YANG, S. L. Further evidence concerning the autoantigenic status of the trophoblast, 1177

—, —, and —. Maternally induced transplantation immunity, tolerance, and runt disease in rats, 808

BENACERRAF, BARUJ. See BLUESTEIN, GREEN, MAURER, and BENACERRAF, 98

—. See WILLIAMS and BENACERRAF, 1279

BERKE, GIDEON, and LEVEY, RAPHAEL H. Cellular immunoabsorbents in transplantation immunity. Specific in vitro deletion and recovery of mouse lymphoid cells sensitized against allogeneic tumors, 972

—, SULLIVAN, KAREN A., and AMOS, BERNARD. Rejection of ascites tumor allografts. I. Isolation, characterization, and in vitro reactivity of peritoneal lymphoid effector cells from BALB/c mice immune to EL4 leukosis, 1334

BERNEY, STEVEN. See ZUCKER-FRANKLIN and BERNEY, 533

BIGGAR, W. D., STUTMAN, OSIAS, and GOOD, ROBERT A. Morphological and functional studies of fetal thymus transplants in mice, 793

BILLINGHAM, R. E. See BEER, BILLINGHAM, and YANG, 808, 1177

BIOZZI, G., STIFFEL, C., MOUTON, D., BOUTHILLIER, Y., and DECREUSEFOND, C. Cytodynamics of the immune response in two lines of mice genetically selected for "high" and "low" antibody synthesis, 1071

BLOMGREN, HENRIC. See GOLSTEIN, WIGZELL, BLOMGREN, and SVEDMYR, 890

BLUESTEIN, HARRY G., GREEN, IRA, MAURER, PAUL H., and BENACERRAF, BARUJ. Specific immune response genes of the guinea pig. V. Influence of the GA and GT immune response genes on the specificity of cellular and humoral immune responses to a terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, 98

BOENISCH, THOMAS. See ALPER, BOENISCH, and WATSON, 68

BOUTHILLIER, Y. See BIOZZI, STIFFEL, MOUTON, BOUTHILLIER, and DECREUSEFOND, 1071

BOYSE, EDWARD A. See SCHEID, BOYSE, CARSWELL, and OLD, 938

BRUNNER, K. T. See SHORTMAN, BRUNNER, and CEROTTINI, 1375

CANADY, MARILYN S. See THOMPSON, PARMELY, FLINK, CANADY, and SEVERSON, 596

CANTOR, HARVEY, and ASOFSKY, RICHARD. Synergy among lymphoid cells mediating the graft-*versus*-host response. III. Evidence for interaction between two types of thymus-derived cells, 764

CARSWELL, ELIZABETH A. See SCHEID, BOYSE, CARSWELL, and OLD, 938

CEROTTINI, J.-C. See SHORTMAN, BRUNNER, and CEROTTINI, 1375

CHAN, EVA L. See MITCHELL, CHAN, NOBLE, WEISSMAN, MISHELL, and HERZENBERG, 165

CHASE, MERRILL W. See MAGUIRE and CHASE, 357

CHOI, YONG SUNG, and GOOD, ROBERT A. Development of chicken lymphoid system. I. Synthesis and secretion of immunoglobulins by chicken lymphoid cells, 1133

COCHRANE, CHARLES G. See WUEPPER and COCHRANE, 1

COHEN, STANLEY. See WARD, COHEN, and FLANAGAN, 1095
COHN, JONATHAN. See ALVARES, LEIGH, COHN, and KAPPAS, 1406
COHN, ZANVIL A. See HOLLAND, HOLLAND, and COHN, 458
—. See WERB and COHN, 21
COLE, GERALD A. See GILDEN, COLE, MONJAN, and NATHANSON, 860
—. See GILDEN, COLE, and NATHANSON, 874
CONE, ROBERT E. See MARCHALONIS, CONE, and ATWELL, 956
COOKE, T. DEREK, HURD, ERIC R., ZIFF, MORRIS, and JASIN, HUGO E. The pathogenesis of chronic inflammation in experimental antigen-induced arthritis. II. Preferential localization of antigen-antibody complexes to collagenous tissues, 323
COONS, ALBERT H. See RUBIN and COONS, 437
COOPER, MAX D. See LAWTON, ASOFSKY, HYLTON, and COOPER, 277
COTTIER, H. See JOEL, HESS, and COTTIER, 907
CUDKOWICZ, GUSTAVO. See MILLER and CUDKOWICZ, 1028
CURRIE, G. A. See GRANT, CURRIE, and ALEXANDER, 150

DAVIE, JOSEPH M., and PAUL, WILLIAM E. Receptors on immunocompetent cells. IV. Direct measurement of avidity of cell receptors and cooperative binding of multivalent ligands, 643
— and —. V. Cellular correlates of the "maturation" of the immune response, 660
DEAK, BEVERLY D. See MCDEVITT, DEAK, SHREFFLER, KLEIN, STIMPFLING, and SNELL, 1259
DECREUSEFOND, C. See BIOZZI, STIFFEL, MOUTON, BOUTHILLIER, and DECREUSEFOND, 1071
DIXON, FRANK J. See LERNER, MCCONAHEY, JANSEN, and DIXON, 136
DIXON, FRANK J. See OLDSTONE and DIXON, 827
DUBOS, RENÉ. See LEE and DUBOS, 220
DUC, H. T. See VOISIN, KINSKY, and DUC, 1185
DVORAK, HAROLD F., and MIHM, MARTIN C., JR. Basophilic leukocytes in allergic contact dermatitis, 235

EIDINGER, D. See HASKILL, ELLIOTT, KERBEL, AXELRAD, and EIDINGER, 1410
ELLIOTT, B. E. See HASKILL, ELLIOTT, KERBEL, AXELRAD, and EIDINGER, 1410
EMESON, EUGENE E. See THURSH and EMESON, 754

FARBER, MIRIAM B. See HUMMELER, HARRIS, HARRIS, and FARBER, 491
FEIZI, TEN, and KABAT, ELVIN A. Immunochemical studies on blood groups. LIV. Classification of anti-I and anti-i sera into groups based on reactivity patterns with various antigens related to the blood group A, B, H, Le^a, Le^b and precursor substances, 1247
FELDMAN, JOSEPH D. See ATKINS, FELDMAN, FRANCIS, and HURSH, 1113
—. See PETER and FELDMAN, 1301
—. See TUBERGEN, FELDMAN, POLLOCK, and LERNER, 255
FELDMANN, MARC. Cell interactions in the immune response in vitro. II. The requirement for macrophages in lymphoid cell collaboration, 1049
—. Induction of immunity and tolerance in vitro by hapten protein conjugates. I. The relationship between the degree of hapten conjugation and the immunogenicity of dinitrophenylated polymerized flagellin, 735

FLANAGAN, THOMAS D. See WARD, COHEN, and FLANAGAN, 1095
FLINK, RONALD J. See THOMPSON, PARMELY, FLINK, CANADY, and SEVERSON, 596
FRANCIS, LORRAINE. See ATKINS, FELDMAN, FRANCIS, and HURSH, 1113
FRIBERG, STEN, JR. See KLEIN, FRIBERG, and HARRIS, 839

GABBIANI, G., HIRSCHEL, B. J., RYAN, G. B., STATKOV, P. R., and MAJNO, G. Granulation tissue as a contractile organ. A study of structure and function, 719
GILDEN, DONALD H., COLE, GERALD A., MONJAN, ANDREW A., and NATHANSON, NEAL. Immunopathogenesis of acute central nervous system disease produced by lymphocytic choriomeningitis virus. I. Cyclophosphamide-mediated induction of the virus-carrier state in adult mice, 860
—, —, and NATHANSON, NEAL. II. Adoptive immunization of virus carriers, 874
GLEICHMANN, ERNST. See GLEICHMANN, GLEICHMANN, ANDRÉ-SCHWARTZ, and SCHWARTZ, 516
GLEICHMANN, HELGA, GLEICHMANN, ERNST, ANDRÉ-SCHWARTZ, JANINE, and SCHWARTZ, ROBERT S. Chronic allogeneic disease. III. Genetic requirements for the induction of glomerulonephritis, 516
GÖTZE, OTTO. See MÜLLER-EBERHARD and GÖTZE, 1003
GOLSTEIN, PIERRE, WIGZELL, HANS, BLOMGREN, HENRIC, and SVEDMYR, ERIK A. J. Cells mediating specific in vitro cytotoxicity. II. Probable autonomy of thymus-processed lymphocytes (T cells) for the killing of allogeneic target cells, 890
GOOD, ROBERT A. See BIGGAR, STUTMAN, and GOOD, 793
—. See CHOI and GOOD, 1133
—. See STUTMAN, YUNIS, and GOOD, 339
GOODMAN, JOEL W. See ALKAN, WILLIAMS, NITECKI, and GOODMAN, 1228
GOWANS, J. L. See HOWARD, HUNT, and GOWANS, 200
GRANT, C. K., CURRIE, G. A., and ALEXANDER, P. Thymocytes from mice immunized against an allograft render bone marrow cells specifically cytotoxic, 150
GREEN, IRA. See BLUESTEIN, GREEN, MAURER, and BENACERRAF, 98
—. See ROSENSTREICH, SHEVACH, GREEN, and ROSENTHAL, 1037
GRUMET, F. CARL. Genetic control of the immune response. A selective defect in immunologic (IgG) memory in nonresponder mice, 110
—. See MITCHELL, GRUMET, and MCDEVITT, 126
GRUNDKE, INGE. See SHERR, BAUR, GRUNDKE, ZELIGS, ZELIGS, and UHR, 1392

HARRIS, HENRY. See KLEIN, FRIBERG, and HARRIS, 839
HARRIS, SUSANNA. See HUMMELER, HARRIS, HARRIS, and FARBER, 491
HARRIS, T. N. See HUMMELER, HARRIS, HARRIS, and FARBER, 491
HART, DAVID A., WANG, AI-LAN, PAWLAK, LAURA L., and NISONOFF, ALFRED. Suppression of idiotypic specificities in adult mice by administration of antiidiotypic antibody, 1293
HASKILL, J. S., ELLIOTT, B. E., KERBEL, R., AXELRAD, M. A., and EIDINGER, D. Classification of thymus-derived and marrow-derived lymphocytes by demonstration of their antigen-binding characteristics, 1410
HAXBY, JAMES A. See KOLB, HAXBY, ARROYAVE, and MÜLLER-EBERHARD, 549
HERZENBERG, L. A. See JACOBSON,

HERZENBERG, RIBLET, and HERZENBERG, 1163
HERZENBERG, L. A. See MITCHELL, CHAN, NOBLE, WEISSMAN, MISHELL, and HERZENBERG, 165
HERZENBERG, LEONORE A. See JACOBSON and HERZENBERG, 1151
—. See JACOBSON, HERZENBERG, RIBLET, and HERZENBERG, 1163
HESS, M. W. See JOEL, HESS, and COTTIER, 907
HIRSCHEL, B. J. See GABBIANI, HIRSCHEL, RYAN, STATKOV, and MAJNO, 719
HOLLAND, NANCY H. See HOLLAND, HOLLAND, and COHN, 458
HOLLAND, PHILLIP, HOLLAND, NANCY H., and COHN, ZANVIL A. The selective inhibition of macrophage phagocytic receptors by anti-membrane antibodies, 458
HOWARD, JONATHAN C. The life-span and recirculation of marrow-derived small lymphocytes from the rat thoracic duct, 185
—, HUNT, S. V., and GOWANS, J. L. Identification of marrow-derived and thymus-derived small lymphocytes in the lymphoid tissue and thoracic duct lymph of normal rats, 200
HUMMELER, KLAUS, HARRIS, T. N., HARRIS, SUSANNA, and FARBER, MIRIAM B. Studies on antibody-producing cells. IV. Ultrastructure of plaque-forming cells of rabbit lymph, 491
HUNT, S. V. See HOWARD, HUNT, and GOWANS, 200
HURD, ERIC R. See COOKE, HURD, ZIFF, and JASIN, 323
HURSH, ELLEN. See ATKINS, FELDMAN, FRANCIS, and HURSH, 1113
HYLTON, MARTHA B. See LAWTON, ASOFSKY, HYLTON, and COOPER, 277

JACOBSON, E. B., and HERZENBERG, LEONORE A. Active suppression of immunoglobulin allotype synthesis. I. Chronic suppression after perinatal exposure to maternal antibody to paternal allotype in (SJL × BALB/c) F_1 mice, 1151
JACOBSON, E. B., HERZENBERG, LEONORE A., RIBLET, R., and HERZENBERG, L. A. Active suppression of immunoglobulin allotype synthesis. II. Transfer of suppressing factor with spleen cells, 1163
JANSEN, INGA. See LERNER, MCCONAHEY, JANSEN, and DIXON, 136
JASIN, HUGO E. See COOKE, HURD, ZIFF, and JASIN, 323
JOEL, D. D., HESS, M. W., and COTTIER, H. Magnitude and pattern of thymic lymphocyte migration in neonatal mice, 907
JOHNSON, ARTHUR G. See STOUT and JOHNSON, 45
JUTILA, JOHN W. See MANNING and JUTILA, 1316

KABAT, ELVIN A. See FEIZI and KABAT, 1247
KAPLAN, ALLEN P., KAY, A. B., and AUSTEN, K. FRANK. A prealbumin activator of prekallikrein. III. Appearance of chemotactic activity for human neutrophils by the conversion of human prekallikrein to kallikrein, 81
KAPPAS, ATTALLAH. See ALVARES, LEIGH, COHN, and KAPPAS, 1406
KARNOVSKY, MORRIS J. See PERKINS, KARNOVSKY, and UNANUE, 267
KAY, A. B. See KAPLAN, KAY, and AUSTEN, 81
KERBEL, R. See HASKILL, ELLIOTT, KERBEL, AXELRAD, and EIDINGER, 1410
KINSKY, R. G. See VOISIN, KINSKY, and DUC, 1185
KLEIN, EVA. See SKURZAK, KLEIN, YOSHIDA, and LAMON, 997
KLEIN, GEORGE, FRIBERG, STEN, JR., and HARRIS, HENRY. Two kinds of

antigen suppression in tumor cells revealed by cell fusion, 839
KLEIN, JAN, and SHREFFLER, DONALD C. Evidence supporting a two-gene model for the H-2 histocompatibility system of the mouse, 924
—. See MCDEVITT, DEAK, SHREFFLER, KLEIN, STIMPFLING, and SNELL, 1259
KOLB, WILLIAM P., HAXBY, JAMES A., ARROYAVE, CARLOS M., and MÜLLER-EBERHARD, HANS J. Molecular analysis of the membrane attack mechanism of complement, 549
KUETTNER, MIRTA GOFFAN, WANG, AI-LAN, and NISONOFF, ALFRED. Quantitative investigations of idiotypic antibodies. VI. Idiotypic specificity as a potential genetic marker for the variable regions of mouse immunoglobulin polypeptide chains, 579

LAFLEUR, LOUIS, MILLER, R. G., and PHILLIPS, R. A. A quantitative assay for the progenitors of bone marrow–associated lymphocytes, 1363
LAMON, EDDIE W. See SKURZAK, KLEIN, YOSHIDA, and LAMON, 997
LANE, F. C., and UNANUE, EMIL R. Requirement of thymus (T) lymphocytes for resistance to listeriosis, 1104
LAWTON, ALEXANDER R. III, ASOFSKY, RICHARD, HYLTON, MARTHA B., and COOPER, MAX D. Suppression of immunoglobulin class synthesis in mice. I. Effects of treatment with antibody to μ-chain, 277
LEE, CHI-JEN, and DUBOS, RENÉ. Lasting biological effects of early environmental influences. VII. Metabolism of adenosine 3′,5′-monophosphate in mice exposed to early environmental stress, 220
LEIGH, SUSAN. See ALVARES, LEIGH, COHN, and KAPPAS, 1406
LERNER, RICHARD A., MCCONAHEY, PATRICIA J., JANSEN, INGA, and DIXON, FRANK J. Synthesis of plasma membrane–associated and secretory immunoglobulin in diploid lymphocytes, 136
—. See TUBERGEN, FELDMAN, POLLOCK, and LERNER, 255
LEROY, E. CARWILE. Connective tissue synthesis by scleroderma skin fibroblasts in cell culture, 1351
LEVEY, RAPHAEL H. See BERKE and LEVEY, 972
LEWIS, HEATHER. See NOSSAL and LEWIS, 1416
—. See NOSSAL, WARNER, LEWIS, and SPRENT, 405
LIEW, F. Y. See PARISH and LIEW, 298

MAGUIRE, HENRY C., JR., and CHASE, MERRILL W. Studies on the sensitization of animals with simple chemical compounds. XIII. Sensitization of guinea pigs with picric acid, 357
MAJNO, G. See GABBIANI, HIRSCHEL, RYAN, STATKOV, and MAJNO, 719
MANDEL, T. See BASTEN, WARNER, and MANDEL, 627
MANNING, DEAN D., and JUTILA, JOHN W. Immunosuppression of mice injected with heterologous anti-immunoglobulin heavy chain antisera, 1316
MANTOVANI, B., RABINOVITCH, M., and NUSSENZWEIG, V. Phagocytosis of immune complexes by macrophages. Different roles of the macrophage receptor sites for complement (C3) and for immunoglobulin (IgG), 780
MARCHALONIS, JOHN J., CONE, ROBERT E., and ATWELL, JOHN L. Isolation and partial characterization of lymphocyte surface immunoglobulins, 956
MARCHESI, VINCENT T. See TILLACK, SCOTT, and MARCHESI, 1209
MAURER, PAUL H. See BLUESTEIN,

GREEN, MAURER, and BENACERRAF, 98
MCCONAHEY, PATRICIA J. See LERNER, MCCONAHEY, JANSEN, and DIXON, 136
MCDEVITT, HUGH O., DEAK, BEVERLY D., SHREFFLER, DONALD C., KLEIN, JAN, STIMPFLING, JACK H., and SNELL, GEORGE D. Genetic control of the immune response. Mapping of the *Ir-1* locus, 1259
—. See MITCHELL, GRUMET, and MCDEVITT, 126
MIHM, MARTIN C., JR. See DVORAK and MIHM, 235
MILLER, HAROLD C., and CUDKOWICZ, GUSTAVO. Immunologic memory cells of bone marrow origin. Increased burst size of specific immunocyte precursors, 1028
MILLER, J. F. A. P. See ANDERSON, SPRENT, and MILLER, 711
—. See BASTEN, MILLER, SPRENT, and PYE, 610
MILLER, R. G. See LAFLEUR, MILLER, and PHILLIPS, 1363
MISHELL, R. I. See MITCHELL, CHAN, NOBLE, WEISSMAN, MISHELL, and HERZENBERG, 165
MITCHELL, GRAHAM F., CHAN, EVA L., NOBLE, MARION S., WEISSMAN, I. L., MISHELL, R. I., and HERZENBERG, L. A. Immunological memory in mice. III. Memory to heterologous erythrocytes in both T cell and B cell populations and requirement for T cells in expression of B cell memory. Evidence using immunoglobulin allotype and mouse alloantigen θ-markers with congenic mice, 165
—, GRUMET, F. CARL, and MCDEVITT, HUGH O. Genetic control of the immune response. The effect of thymectomy on the primary and secondary antibody response of mice to poly-L (Tyr, Glu)-poly-D, L-Ala--poly-L-Lys, 126
MONJAN, ANDREW A. See GILDEN, COLE, MONJAN, and NATHANSON, 860
MOUTON, D. See BIOZZI, STIFFEL, MOUTON, BOUTHILLIER, and DECREUSEFOND, 1071
MOZES, EDNA. See SHEARER, MOZES, and SELA, 1009
MÜLLER-EBERHARD, HANS J., and GÖTZE, OTTO. C3 proactivator convertase and its mode of action, 1003
—. See KOLB, HAXBY, ARROYAVE, and MÜLLER-EBERHARD, 549

NAKAMURA, KUNIE. The proliferation of plasma cells from mouse bone marrow in vitro. I. The role of thymus, 476
NATHANSON, NEAL. See GILDEN, COLE, MONJAN, and NATHANSON, 860
—. See GILDEN, COLE, and NATHANSON, 874
NISHIO, J. See SALVIN and NISHIO, 985
NISONOFF, ALFRED. See HART, WANG, PAWLAK, and NISONOFF, 1293
—. See KUETTNER, WANG, and NISONOFF, 579
NITECKI, DANUTE E. See ALKAN, WILLIAMS, NITECKI, and GOODMAN, 1228
NOBLE, MARION S. See MITCHELL, CHAN, NOBLE, WEISSMAN, MISHELL, and HERZENBERG, 165
NOSSAL, G. J. V., and LEWIS, HEATHER. Variation in accessible cell surface immunoglobulin among antibody-forming cells, 1416
—, WARNER, N. L., LEWIS, HEATHER, and SPRENT, J. Quantitative features of a sandwich radioimmunolabeling technique for lymphocyte surface receptors, 405
NUSSENZWEIG, V. See MANTOVANI, RABINOVITCH, and NUSSENZWEIG, 780

OLD, LLOYD J. See SCHEID, BOYSE, CARSWELL, and OLD, 938
OLDSTONE, MICHAEL B. A., and

DIXON, FRANK J. Disease accompanying *in utero* viral infection. The role of maternal antibody in tissue injury after transplacental infection with lymphocytic choriomeningitis virus, 827

PARISH, C. R., and LIEW, F. Y. Immune response to chemically modified flagellin. III. Enhanced cell-mediated immunity during high and low zone antibody tolerance to flagellin, 298

PARMELY, MICHAEL J. See THOMPSON, PARMELY, FLINK, CANADY, and SEVERSON, 596

PAUL, WILLIAM E. See DAVIE and PAUL, 643, 660

PAWLAK, LAURA L. See HART, WANG, PAWLAK, and NISONOFF, 1293

PERKINS, WILLIAM D., KARNOVSKY, MORRIS J., and UNANUE, EMIL R. An ultrastructural study of lymphocytes with surface-bound immunoglobulin, 267

PETER, HANS-HARTMUT, and FELDMAN, JOSEPH D. Cell-mediated cytotoxicity during rejection and enhancement of allogeneic skin grafts in rats, 1301

PHILLIPS, R. A. See LAFLEUR, MILLER, and PHILLIPS, 1363

PIERCE, CARL W., SOLLIDAY, SUSAN M., and ASOFSKY, RICHARD. Immune responses in vitro. IV. Suppression of primary γM, γG, and γA plaque-forming cell responses in mouse spleen cell cultures by class-specific antibody to mouse immunoglobulins, 675

—, —, and —. V. Suppression of γM, γG, and γA plaque-forming cell responses in cultures of primed mouse spleen cells by class-specific antibody to mouse immunoglobulins, 698

PINCUS, THEODORE. See ROWE and PINCUS, 429

POLLOCK, E. M. See TUBERGEN, FELDMAN, POLLOCK, and LERNER, 255

PYE, J. See BASTEN, MILLER, SPRENT, and PYE, 610

RABINOVITCH, M. See MANTOVANI, RABINOVITCH, and NUSSENZWEIG, 780

RIBLET, R. See JACOBSON, HERZENBERG, RIBLET, and HERZENBERG, 1163

ROSENSTREICH, DAVID L., SHEVACH, ETHAN, GREEN, IRA, and ROSENTHAL, ALAN S. The uropod-bearing lymphocyte of the guinea pig. Evidence for thymic origin, 1037

ROSENTHAL, ALAN S. See ROSENSTREICH, SHEVACH, GREEN, and ROSENTHAL, 1037

ROWE, WALLACE P., and PINCUS, THEODORE. Quantitative studies of naturally occurring murine leukemia virus infection of AKR mice, 429

RUBIN, ARNOLD S., and COONS, ALBERT H. Specific heterologous enhancement of immune responses. II. Immunological memory cells of thymic origin, 437

RYAN, G. B. See GABBIANI, HIRSCHEL, RYAN, STATKOV, and MAJNO, 719

SALVIN, S. B., and NISHIO, J. Lymphoid cells in delayed hypersensitivity. III. The influence of X-irradiation on passive transfer and on in vitro production of soluble mediators, 985

SCHEID, MARGRIT, BOYSE, EDWARD A., CARSWELL, ELIZABETH A., and OLD, LLOYD J. Serologically demonstrable alloantigens of mouse epidermal cells, 938

SCHWARTZ, ROBERT S. See GLEICHMANN, GLEICHMANN, ANDRÉ-SCHWARTZ, and SCHWARTZ, 516

SCOTT, ROBERT E. See TILLACK, SCOTT, and MARCHESI, 1209

SELA, MICHAEL. See SHEARER, MOZES, and SELA, 1009

SEVERSON, CHARLES D. See THOMP-

SON, PARMELY, FLINK, CANADY, and SEVERSON, 596
SHEARER, G. M., MOZES, EDNA, and SELA, MICHAEL. Contribution of different cell types to the genetic control of immune responses as a function of the chemical nature of the polymeric side chains (poly-L-prolyl and poly-DL-alanyl) of synthetic immunogens, 1009
SHERR, CHARLES J., BAUR, SONIA, GRUNDKE, INGE, ZELIGS, JOSEPH, ZELIGS, BARBARA, and UHR, JONATHAN W. Cell surface immunoglobulin. III. Isolation and characterization of immunoglobulin from nonsecretory human lymphoid cells, 1392
SHEVACH, ETHAN. See ROSENSTREICH, SHEVACH, GREEN, and ROSENTHAL, 1037
SHORTMAN, K., BRUNNER, K. T., and CEROTTINI, J.-C. Separation of stages in the development of the "T" cells involved in cell-mediated immunity, 1375
SHREFFLER, DONALD C. See KLEIN and SHREFFLER, 924
—. See MCDEVITT, DEAK, SHREFFLER, KLEIN, STIMPFLING, and SNELL, 1259
SILVERS, WILLYS K. See WACHTEL and SILVERS, 388
SJÖBERG, OLOF. Rapid breaking of tolerance against *Escherichia coli* lipopolysaccharide in vivo and in vitro, 850
SKURZAK, HENRYK M., KLEIN, EVA, YOSHIDA, TAKATO O., and LAMON, EDDIE W. Synergistic or antagonistic effect of different antibody concentrations on in vitro lymphocyte cytotoxicity in the Moloney sarcoma virus system, 997
SNELL, GEORGE D. See MCDEVITT, DEAK, SHREFFLER, KLEIN, STIMPFLING, and SNELL, 1259
SOLLIDAY, SUSAN M. See PIERCE, SOLLIDAY, and ASOFSKY, 675, 698
SPRENT, J. See ANDERSON, SPRENT, and MILLER, 711
SPRENT, J. See BASTEN, MILLER, SPRENT, and PYE, 610
—. See NOSSAL, WARNER, LEWIS, and SPRENT, 405
STATKOV, P. R. See GABBIANI, HIRSCHEL, RYAN, STATKOV, and MAJNO, 719
STEPHENSON, JOHN R., and AARONSON, STUART A. Antigenic properties of murine sarcoma virus–transformed BALB/3T3 nonproducer cells, 503
STIFFEL, C. See BIOZZI, STIFFEL, MOUTON, BOUTHILLIER, and DECREUSEFOND, 1071
STIMPFLING, JACK H. See MCDEVITT, DEAK, SHREFFLER, KLEIN, STIMPFLING, and SNELL, 1259
STOUT, ROBERT D., and JOHNSON, ARTHUR G. Regulation of the immune system by synthetic polynucleotides. V. Effect on cell-associated immunoglobulin receptors and immunological memory, 45
STREILEIN, J. WAYNE. Analysis of graft-*versus*-host disease in Syrian hamsters. IV. The refractory state and immunologic competence, 567
STUTMAN, OSIAS, YUNIS, EDMOND J., and GOOD, ROBERT A. Studies on thymus function. III. Duration of thymic function, 339
—. See BIGGAR, STUTMAN, and GOOD, 793
SULLIVAN, KAREN A. See BERKE, SULLIVAN, and AMOS, 1334
SVEDMYR, ERIK A. J. See GOLSTEIN, WIGZELL, BLOMGREN, and SVEDMYR, 890

TAKAHASHI, TOSHITADA. See AOKI and TAKAHASHI, 443
THOMPSON, JOHN S., PARMELY, MICHAEL J., FLINK, RONALD J., CANADY, MARILYN S., and SEVERSON, CHARLES D. Relationship of mixed lymphocyte culture response to *HL-A* histocompatibility antigens. Effect of allele plus one antigen match, 596
THURSH, DONALD R., and EMESON,

EUGENE E. The immunologically specific retention of recirculating long-lived lymphocytes in lymph nodes stimulated by xenogeneic erythrocytes, 754

TIGELAAR, ROBERT E., and ASOFSKY, RICHARD. Synergy among lymphoid cells mediating the graft-*versus*-host response. IV. Synergy in the GVH reaction quantitated by a mortality assay in sublethally irradiated recipients, 1059

TILLACK, THOMAS W., SCOTT, ROBERT E., and MARCHESI, VINCENT T. The structure of erythrocyte membranes studied by freeze-etching. II. Localization of receptors for phytohemagglutinin and influenza virus to the intramembranous particles, 1209

TUBERGEN, DAVID G., FELDMAN, JOSEPH D., POLLOCK, E. M., and LERNER, RICHARD A. Production of macrophage migration inhibition factor by continuous cell lines, 255

UHR, JONATHAN W. See SHERR, BAUR, GRUNDKE, ZELIGS, ZELIGS, and UHR, 1392

UNANUE, EMIL R. See LANE and UNANUE, 1104

—. See PERKINS, KARNOVSKY, and UNANUE, 267

VOISIN, G. A., KINSKY, R. G., and DUC, H. T. Immune status of mice tolerant of living cells. II. Continuous presence and nature of facilitation-enhancing antibodies in tolerant animals, 1185

WACHTEL, STEPHEN S., and SILVERS, WILLYS K. The role of passenger leukocytes in the anomalous survival of neonatal skin grafts in mice, 388

WANG, AI-LAN. See HART, WANG, PAWLAK, and NISONOFF, 1293

—. See KUETTNER, WANG, and NISONOFF, 579

WARD, PETER A., COHEN, STANLEY, and FLANAGAN, THOMAS D. Leukotactic factors elaborated by virus-infected tissues, 1095

WARNER, N. L. See BASTEN, WARNER, and MANDEL, 627

—. See NOSSAL, WARNER, LEWIS, and SPRENT, 405

WATSON, LILLIAN. See ALPER, BOENISCH, and WATSON, 68

WEISSMAN, I. L. See MITCHELL, CHAN, NOBLE, WEISSMAN, MISHELL, and HERZENBERG, 165

WERB, ZENA, and COHN, ZANVIL A. Cholesterol metabolism in the macrophage. III. Ingestion and intracellular fate of cholesterol and cholesterol esters, 21

WIDMER, M. B., and BACH, F. H. Allogeneic and xenogeneic response in mixed leukocyte cultures, 1204

WIGZELL, HANS. See GOLSTEIN, WIGZELL, BLOMGREN, and SVEDMYR, 890

WILLIAMS, E. BRADY. See ALKAN, WILLIAMS, NITECKI, and GOODMAN, 1228

WILLIAMS, R. MICHAEL, and BENACERRAF, BARUJ. Genetic control of thymus-derived cell function. I. In vitro DNA synthetic response of normal mouse spleen cells stimulated by the mitogens concanavalin A and phytohemagglutinin, 1279

WUEPPER, KIRK D., and COCHRANE, CHARLES G. Plasma prekallikrein: isolation, characterization, and mechanism of activation, 1

YANG, S. L. See BEER, BILLINGHAM, and YANG, 808, 1177

YOSHIDA, TAKATO O. See SKURZAK,

KLEIN, YOSHIDA, and LAMON, 997
YUNIS, EDMOND J. See STUTMAN, YUNIS, and GOOD, 339

ZELIGS, BARBARA. See SHERR, BAUR, GRUNDKE, ZELIGS, ZELIGS, and UHR, 1392
ZELIGS, JOSEPH. See SHERR, BAUR, GRUNDKE, ZELIGS, ZELIGS, and UHR, 1392
ZIFF, MORRIS. See COOKE, HURD, ZIFF, and JASIN, 323
ZUCKER-FRANKLIN, DOROTHEA, and BERNEY, STEVEN. Electron microscope study of surface immunoglobulin–bearing human tonsil cells, 533

INDEX TO SUBJECTS

ACID, deoxyribonucleic, response, synthetic in vitro, of spleen cells stimulated by mitogens concanavalin A and phytohemagglutinin, 1279

—, L-glutamic, L-alanine, and L-tyrosine, terpolymer of, influence of GA and GT immune response genes on specificity of cellular and humoral immune responses to, 98

—, picric, sensitization with simple chemical compounds, 357

Adenosine 3′,5′-monophosphate, metabolism in mice exposed to early environmental stress, 220

L-Alanine, L-glutamic acid, and L-tyrosine, terpolymer of, influence of GA and GT immune response genes on specificity of cellular and humoral immune responses to, 98

Allergic contact dermatitis, basophilic leukocytes in, 235

Alloantigen, mouse, θ-markers with congenic mice, evidence for immunological memory to heterologous erythrocytes in both T and B cell populations and requirement for T cells in expression of B cell memory using, 165

Alloantigens of epidermal cells, serologically demonstrable, 938

Allogeneic disease, chronic, genetic requirements for induction of glomerulonephritis, 516

— skin grafts, cell-mediated cytotoxicity during rejection and enhancement of, 1301

— target cells, probable autonomy of thymus-processed lymphocytes, T cells, for killing of, 890

— tumors, specific in vitro deletion and recovery of lymphoid cells sensitized against, 972

Allogeneic and xenogeneic response in mixed leukocyte cultures, 1204

Allograft, thymocytes from mice immunized against render bone marrow cells specifically cytotoxic, 150

Allografts, tumor, ascites, rejection, 1334

Allotype, immunoglobulin, and mouse alloantigen θ-markers with congenic mice, evidence for immunological memory to heterologous erythrocytes in both T and B cell populations and requirement for T cells in expression of B cell memory using, 165

— synthesis, immunoglobulin, active suppression of, 1151, 1163

Antibodies, anti-membrane, selective inhibition of macrophage phagocytic receptors by, 458

—, facilitation-enhancing, continuous presence and nature in tolerant animals, 1185

—, idiotypic specificity as potential genetic marker for variable regions of immunoglobulin polypeptide chains, 579

Antibody, antiidiotypic, suppression of idiotypic specificities by administration of, 1293

—, class-specific, to immunoglobulins, suppression of γM, γG, and γA plaque-forming cell responses in cultures of primed spleen cells by, 698

—, class-specific, to immunoglobulins, suppression of primary γM, γG, and γA plaque-forming cell responses in spleen cell cultures by, 675

Antibody concentrations, different, synergistic or antagonistic effect on in vitro lymphocyte cytotoxicity in Moloney sarcoma virus system, 997
—, maternal, role in tissue injury after transplacental infection with lymphocytic choriomeningitis virus, 827
—, maternal, to paternal allotype in (SJL × BALB/c)F_1 mice, chronic suppression after perinatal exposure to, 1151
— receptor on B lymphocytes, detection method and functional significance, 610
— receptor on B lymphocytes, immunochemical and electron microscopy characteristics, 627
— response, primary and secondary, to poly-L(Tyr,Glu)-poly-D,L-Ala–poly-L-Lys, effect of thymectomy on, 126
—, secreted, and cellular receptor for antigen on immunological memory cells, correlation between binding properties, 312
— synthesis, high and low, cytodynamics of immune response in two lines of mice genetically selected for, 1071
— to μ-chain, effects of treatment with on suppression of immunoglobulin class synthesis, 277
— tolerance, high and low zone, to flagellin, enhanced cell-mediated immunity during, 298
Antibody-antigen complexes, preferential localization to collagenous tissues in antigen-induced arthritis, 323
Antibody-forming cells, variation in accessible cell surface immunoglobulin among, 1416
Antibody-producing cells, ultrastructure of plaque-forming cells of lymph, 491
Antigen, cellular receptor on immunological memory cells and secreted antibody, correlation between binding properties, 312
— molecules, small mono- and bifunctional, humoral and cellular immune responses to, 1228
Antigen recognition and immune response, 1228
— suppression, two kinds, in tumor cells, revealed by cell fusion, 839
Antigen-antibody complexes, preferential localization to collagenous tissues in antigen-induced arthritis, 323
Antigen-binding characteristics, classification of thymus-derived and marrow-derived lymphocytes by demonstration of, 1410
Antigenic properties of murine sarcoma virus–transformed BALB/3T3 nonproducer cells, 503
Antigen-induced arthritis, pathogenesis of chronic inflammation in, 323
Antigens, histocompatibility, *HL-A*, relationship of mixed lymphocyte culture response to, effect of allele plus one antigen match, 596
—, surface, viral and cellular, of murine leukemias and myelomas, serological analysis by immunoelectron microscopy, 443
—, various, related to blood group A, B, H, Le^a, Le^b and precursor substances, classification of anti-I and anti-i sera into groups based on reactivity patterns with, 1247
Antisera, heavy chain, anti-immunoglobulin, heterologous, immunosuppression of mice injected with, 1316
Arthritis, antigen-induced, pathogenesis of chronic inflammation in, 323
Ascites tumor allografts, rejection, 1334
Autoantigenic status of trophoblast, further evidence concerning, 1177

BACTERIAL chemotactic factor and chemotactic behavior of complement-derived factors, C3a, C5a, and $C\overline{567}$, relationship to

ability to activate proesterase 1 of polymorphonuclear leukocytes, 376
Basophilic leukocytes in allergic contact dermatitis, 235
Biological effects, lasting, of early environmental influences, 220
Blood group A, B, H, Le^a, Le^b and precursor substances, classification of anti-I and anti-i sera into groups based on reactivity patterns with various antigens related to, 1247
Bone marrow cells, thymocytes from mice immunized against allograft render specifically cytotoxic, 150
— marrow, in vitro proliferation of plasma cells from, role of thymus, 476
— marrow origin, immunologic memory cells of, 1028

CELL collaboration, lymphoid, requirement for macrophages in, 1049
— culture, connective tissue synthesis by scleroderma skin fibroblasts in, 1351
— fusion, two kinds of antigen suppression in tumor cells revealed by, 839
— interactions in immune response in vitro, 1049
— lines, continuous, production of macrophage migration inhibition factor by, 255
— population, both T and B, immunological memory to heterologous erythrocytes in, and requirement for T cells in expression of B cell memory, evidence using immunoglobulin allotype and mouse alloantigen θ-marker with congenic mice, 165
— receptors, direct measurement of avidity and cooperative binding of multivalent ligands, 643
— responses, plaque-forming, γM, γG, and γA, suppression in cultures of primed spleen cells by class-specific antibody to immunoglobulins, 698
Cell responses, plaque-forming, primary γM, γG, and γA, suppression in spleen cell cultures by class-specific antibody to immunoglobulins, 675
— surface immunoglobulin, accessible, variation among antibody-forming cells, 1416
— surface immunoglobulin, isolation and characterization from nonsecretory human lymphoid cells, 1392
—, thymus-derived, genetic control of function, 1279
— types, different, contribution to genetic control of immune responses as function of chemical nature of polymeric side chains (poly-L-prolyl and poly-DL-alanyl) of synthetic immunogens, 1009
Cell-associated immunoglobulin receptors and immunological memory, effect of synthetic polynucleotides on, 45
Cell-mediated cytotoxicity during rejection and enhancement of allogeneic skin grafts, 1301
— immunity, enhanced, during high and low zone antibody tolerance to flagellin, 298
— immunity, separation of stages in development of T cells involved in, 1375
Cells, antibody-forming, variation in accessible cell surface immunoglobulin among, 1416
—, antibody-producing, ultrastructure of plaque-forming cells of lymph, 491
—, bone marrow, thymocytes from mice immunized against allograft render specifically cytotoxic, 150
—, epidermal, serologically demonstrable alloantigens of, 938
—, immunocompetent, receptors on, cellular correlates of maturation of immune response, 660
—, immunocompetent, receptors on, direct measurement of avidity and cooperative binding of multivalent ligands, 643

Cells, immunologic memory, of bone marrow origin, 1028
—, immunological memory, of thymic origin, 437
—, living, immune status of mice tolerant of, 1185
—, lymphoid, chicken, synthesis and secretion of immunoglobulins by, 1133
—, lymphoid effector, peritoneal, from BALB/c mice immune to EL4 leukosis, isolation, characterization, and in vitro reactivity of, 1334
—, lymphoid, human, nonsecretory, isolation and characterization of cell surface immunoglobulin from, 1392
—, lymphoid, in delayed hypersensitivity, 985
—, lymphoid, mediating graft-*versus*-host response, synergy among, 764, 1059
—, lymphoid, sensitized against allogeneic tumors, specific in vitro deletion and recovery of, 972
— mediating specific in vitro cytotoxicity, 890
—, memory, immunological, correlation between binding properties of secreted antibody and cellular receptor for antigen on, 312
—, murine sarcoma virus–transformed BALB/3T3 nonproducer, antigenic properties of, 503
—, plasma, proliferation from bone marrow in vitro, role of thymus, 476
—, primed spleen, cultures, suppression of γM, γG, and γA plaque-forming cell responses in by class-specific antibody to immunoglobulins, 698
—, spleen, stimulated by mitogens concanavalin A and phytohemagglutinin, in vitro DNA synthetic response of, 1279
—, spleen, transfer of suppressing factor with, active suppression of immunoglobulin allotype synthesis, 1163
—, thymus-derived, two types, evidence for interaction between in graft-*versus*-host response, 764
Cells, T, involved in cell-mediated immunity, separation of stages in development of, 1375
—, T, thymus-processed lymphocytes, probable autonomy for killing of allogeneic target cells, 890
—, tonsil, human, surface immunoglobulin–bearing, electron microscope study, 533
—, tumor, antigen suppression in, two kinds revealed by cell fusion, 839
Cell-to-cell interaction in immune response, radiosensitivity of thymus-derived lymphocytes, 711
Cellular correlates of maturation of immune response, 660
— and humoral immune responses to small mono- and bifunctional antigen molecules, 1228
— and humoral immune responses to terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, influence of GA and GT immune response genes on specificity of, 98
— immunoabsorbents in transplantation immunity, 972
— receptor for antigen on immunological memory cells, and secreted antibody, correlation between binding properties, 312
— and viral surface antigens of murine leukemias and myelomas, serological analysis by immunoelectron microscopy, 443
μ-Chain, antibody to, effects of treatment with on suppression of immunoglobulin class synthesis, 277
Chain, heavy, antisera, heterologous anti-immunoglobulin, immunosuppression of mice injected with, 1316
Chains, polymeric side (poly-L-prolyl and poly-DL-alanyl), of synthetic immunogens, contribution of different cell types to genetic control of immune responses as function of chemical nature of, 1009
—, polypeptide, immunoglobulin, idiotypic specificity as potential

genetic marker for variable regions of, 579
Chemical compounds, simple, picric acid, sensitization with, 357
—, immuno-, and electron microscopy characteristics of antibody receptor on B lymphocytes, 627
—, immuno-, studies on blood groups, 1247
— nature of polymeric side chains (poly-L-prolyl and poly-DL-alanyl) of synthetic immunogens, contribution of different cell types to genetic control of immune responses as function of, 1009
Chemically modified flagellin, immune response to, 298
Chemotactic activity for human neutrophils by conversion of human prekallikrein to kallikrein, appearance of, 81
— behavior of complement-derived factors, C3a, C5a, and C$\overline{567}$, and bacterial chemotactic factor, relationship to ability to activate proesterase 1 of polymorphonuclear leukocytes, 376
Cholesterol metabolism in macrophage, 21
Choriomeningitis virus, lymphocytic, immunopathogenesis of acute central nervous system disease produced by, 860, 874
— virus, lymphocytic, role of maternal antibody in tissue injury after transplacental infection with, 827
Collagenous tissues, preferential localization of antigen-antibody complexes to in antigen-induced arthritis, 323
Complement (C3) and immunoglobulin (IgG), different roles of macrophage receptor sites for, 780
—, membrane attack mechanism, molecular analysis, 549
Complement-derived factors, C3a, C5a, and C$\overline{567}$, chemotactic behavior, and bacterial chemotactic factor, relationship to ability to activate proesterase 1 of polymorphonuclear leukocytes, 376
Concanavalin A and phytohemagglutinin, mitogens, in vitro DNA synthetic response of spleen cells stimulated by, 1279
Congenic mice, memory to heterologous erythrocytes in both T and B cell populations and requirement for T cells in expression of B cell memory, evidence using immunoglobulin allotype and mouse alloantigen θ-markers with, 165
Conjugates, hapten protein, induction of immunity and tolerance in vitro by, 735
Conjugation, hapten, degree, and immunogenicity of dinitrophenylated polymerized flagellin, relationship between, 735
Connective tissue synthesis by scleroderma skin fibroblasts in cell culture, 1351
Contractile organ, granulation tissue as, structure and function, 719
Convertase, C3 proactivator and its mode of action, 1003
Culture, cell, connective tissue synthesis by scleroderma skin fibroblasts in, 1351
—, mixed, lymphocyte, relationship of response to *HL-A* histocompatibility antigens, effect of allele plus one antigen match, 596
Cultures, leukocyte, mixed, allogeneic and xenogeneic response in, 1204
—, primed spleen cells, suppression of γM, γG, and γA plaque-forming cell responses in by class-specific antibody to immunoglobulins, 698
—, spleen cell, suppression of primary γM, γG, and γA plaque-forming cell responses in by class-specific antibody to immunoglobulins, 675
Cyclophosphamide-mediated induc-

tion of virus-carrier state in immunopathogenesis of acute central nervous system disease produced by lymphocytic choriomeningitis virus, 860
Cytochrome P-450 and mixed function oxidase system in liver, effects of acute exposure to lead and methyl mercury on, 1406
Cytodynamics of immune response in two lines of mice genetically selected for high and low antibody synthesis, 1071
Cytotoxic, specifically, thymocytes from mice immunized against allograft render bone marrow cells, 150
Cytotoxicity, cell-mediated, during rejection and enhancement of allogeneic skin grafts, 1301
—, in vitro, specific, cells mediating, 890
—, lymphocyte, in vitro in Moloney sarcoma virus system, synergistic or antagonistic effect of different antibody concentrations on, 997

DEOXYRIBONUCLEIC acid response, synthetic in vitro, of spleen cells stimulated by mitogens concanavalin A and phytohemagglutinin, 1279
Dermatitis, allergic contact, basophilic leukocytes in, 235
Dinitrophenylated polymerized flagellin, immunogenicity, and degree of hapten conjugation, relationship between, 735
Disease accompanying *in utero* viral infection, role of maternal antibody in tissue injury after transplacental infection with lymphocytic choriomeningitis virus, 827
—, allogeneic, chronic, genetic requirements for induction of glomerulonephritis, 516
—, central nervous system, acute, produced by lymphocytic choriomeningitis virus, immunopathogenesis, 860, 874
—, graft-*versus*-host, in Syrian hamsters, refractory state and immunologic competence, 567
Disease, runt, maternally induced transplantation immunity, and tolerance in rats, 808
Duct, lymph, thoracic, and lymphoid tissue, identification of marrow-derived and thymus-derived small lymphocytes in, 200
—, thoracic, life-span and recirculation of marrow-derived small lymphocytes from, 185

ELECTRON, immuno-, microscopy, serological analysis of viral and cellular surface antigens of murine leukemias and myelomas by, 443
— microscope study of surface immunoglobulin–bearing human tonsil cells, 533
— microscopy and immunochemical characteristics of antibody receptor on B lymphocytes, 627
Environmental influences, early, lasting biological effects, 220
Epidermal cells, serologically demonstrable alloantigens of, 938
Erythrocyte membranes, structure studied by freeze-etching, 1209
Erythrocytes, heterologous, in both T and B cell populations, immunological memory to, and requirement for T cells in expression of B cell memory, evidence using immunoglobulin allotype and mouse alloantigen θ-markers with congenic mice, 165
—, xenogeneic, immunologically specific retention of recirculating long-lived lymphocytes in lymph nodes stimulated by, 754
Escherichia coli lipopolysaccharide, tolerance against, rapid breaking in vivo and in vitro, 850
Esters, cholesterol, and cholesterol, ingestion and intracellular fate in macrophage, 21

FETAL thymus transplants, morphological and functional studies, 793

Fever accompanying delayed hypersensitivity, mechanism, 1113
Fibroblasts, skin, scleroderma, in cell culture, connective tissue synthesis by, 1351
Flagellin, dinitrophenylated polymerized, immunogenicity, and degree of hapten conjugation, relationship between, 735
—, enhanced cell-mediated immunity during high and low zone antibody tolerance to, 298
Freeze-etching, structure of erythrocyte membranes studied by, 1209

GENE, two-, model for H-Z histocompatibility system of mouse, evidence supporting, 924
Genes, specific immune response, GA and GT, influence on specificity of cellular and humoral immune responses to terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, 98
Genetic control of immune response, 126
— control of immune response, mapping of *Ir-1* locus, 1259
— control of immune response, selective defect in immunologic (IgG) memory in nonresponder mice, 110
— control of immune responses as function of chemical nature of polymeric side chains (poly-L-prolyl and poly-DL-alanyl) of synthetic immunogens, contribution of different cell types to, 1009
— control of thymus-derived cell function, 1279
— marker, potential, for variable regions of immunoglobulin polypeptide chains, idiotypic specificity as, 579
— polymorphism in human glycine-rich β-glycoprotein, 68
— requirements for induction of glomerulonephritis in chronic allogeneic disease, 516
— selection for high and low antibody synthesis in two lines of mice, cytodynamics of immune response in, 1071
Glomerulonephritis, genetic requirements for induction in chronic allogeneic disease, 516
L-Glutamic acid, L-alanine, and L-tyrosine, terpolymer of, influence of GA and GT immune response genes on specificity of cellular and humoral immune responses to, 98
Glycine-rich β-glycoprotein, human, genetic polymorphism in, 68
β-Glycoprotein, glycine-rich, human, genetic polymorphism in, 68
Graft-*versus*-host disease in Syrian hamsters, refractory state and immunologic competence, 567
— response, synergy among lymphoid cells mediating, 764, 1059
—, skin, neonatal, role of passenger leukocytes in anomalous survival of, 388
Grafts, skin, allogeneic, cell-mediated cytotoxicity during rejection and enhancement of, 1301
Granulation tissue as contractile organ, structure and function, 719

HAPTEN protein conjugation, degree, and immunogenicity of dinitrophenylated polymerized flagellin, relationship between, 735
Heterologous anti-immunoglobulin heavy chain antisera, immunosuppression of mice injected with, 1316
— enhancement, specific, of immune responses, 437
— erythrocytes in both T and B cell populations, immunological memory to, and requirement for T cells in expression of B cell memory, evidence using immunoglobulin allotype and mouse alloantigen θ-markers with congenic mice, 165
Histocompatibility antigens, *HL-A*, relationship of mixed lymphocyte culture response to, effect of allele plus one antigen match, 596
— system, H-2, of mouse, evidence supporting two-gene model for, 924

Host-*versus*-graft disease in Syrian hamsters, refractory state and immunologic competence, 567
— response, synergy among lymphoid cells mediating, 764, 1059
Humoral and cellular immune responses to small mono- and bifunctional antigen molecules, 1228
— and cellular immune responses to terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, influence of GA and GT immune response genes on specificity of, 98
Hypersensitivity, delayed, lymphoid cells in, 985
—, delayed, mechanism of fever accompanying, 1113

IDIOTYPIC specificities, suppression by administration of anti-idiotypic antibody, 1293
— specificity as potential genetic marker for variable regions of immunoglobulin polypeptide chains, 579
Immune complexes, phagocytosis by macrophages, 780
— response, cell-to-cell interaction in, radiosensitivity of thymus-derived lymphocytes, 711
— response, cellular correlates of maturation of, 660
— response genes, specific, GA and GT, influence on specificity of cellular and humoral immune responses to terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, 98
— response, genetic control, 126
— response, genetic control, mapping of *Ir-1* locus, 1259
— response, genetic control, selective defect in immunologic (IgG) memory in nonresponder mice, 110
— response in two lines of mice genetically selected for high and low antibody synthesis, cytodynamics of, 1071
— response in vitro, cell interactions in, 1049
— response to chemically modified flagellin, 298
— responses as function of chemical nature of polymeric side chains (poly-L-prolyl and poly-DL-alanyl) of synthetic immunogens, contribution of different cell types to genetic control of, 1009
Immune responses, humoral and cellular, to small mono- and bifunctional antigen molecules, 1228
— responses in vitro, suppression of γM, γG, and γA plaque-forming cell responses in cultures of primed spleen cells by class-specific antibody to immunoglobulins, 698
— responses in vitro, suppression of primary γM, γG, and γA plaque-forming cell responses in spleen cell cultures by class-specific antibody to immunoglobulins, 675
— responses, specific heterologous enhancement of, 437
— status of mice tolerant of living cells, 1185
— system, regulation by synthetic polynucleotides, 45
Immunity, cell-mediated, enhanced, during high and low zone antibody tolerance to flagellin, 298
—, cell-mediated, separation of stages in development of T cells involved in, 1375
— to EL4 leukosis, BALB/c mice, isolation, characterization, and in vitro reactivity of peritoneal lymphoid effector cells from, 1334
— and tolerance, induction in vitro by hapten protein conjugates, 735
—, transplantation, cellular, immunoabsorbents in, 972
—, transplantation, maternally induced, tolerance, and runt disease in rats, 808
Immunization, adoptive, of virus carriers in immunopathogenesis of acute central nervous system disease produced by lymphocytic choriomeningitis virus, 874
— of thymocytes from mice against allograft render bone marrow cells specifically cytotoxic, 150
Immunoabsorbents, cellular, in transplantation immunity, 972

Immunochemical and electron microscopy characteristics of antibody receptor on B lymphocytes, 627
— studies on blood groups, 1247
Immunocompetent cells, receptors on, cellular correlates of maturation of immune response, 660
— cells, receptors on, direct measurement of avidity and cooperative binding of multivalent ligands, 643
Immunocyte precursors, specific, increased burst size of, 1028
Immunoelectron microscopy, serological analysis of viral and cellular surface antigens of murine leukemias and myelomas by, 443
Immunogenicity of dinitrophenylated polymerized flagellin and degree of hapten conjugation, relationship between, 735
Immunogens, synthetic, polymeric side chains (poly-L-prolyl and poly-DL-alanyl), contribution of different cell types to genetic control of immune responses as function of chemical nature of, 1009
Immunoglobulin allotype and mouse alloantigen θ-markers with congenic mice, evidence for immunological memory to heterologous erythrocytes in both T and B cell populations and requirement for T cells in expression of B cell memory using, 165
— allotype synthesis, active suppression of, 1151, 1163
—, anti-, heavy chain antisera, heterologous, immunosuppression of mice injected with, 1316
—, cell surface, accessible, variation among antibody-forming cells, 1416
—, cell surface, isolation and characterization from nonsecretory human lymphoid cells, 1392
—, class synthesis, suppression, effects of treatment with antibody to μ-chain, 277
— (IgG) and complement (C3), different roles of macrophage receptor sites for, 780
Immunoglobulin, plasma membrane-associated and secretory, synthesis in diploid lymphocytes, 136
— polypeptide chains, idiotypic specificity as potential genetic marker for variable regions of, 579
— receptors, cell-associated, and immunological memory, effect of synthetic polynucleotides on, 45
—, surface-bound, ultrastructural study of lymphocytes with, 267
Immunoglobulin-bearing human tonsil cells, surface, electron microscope study, 533
Immunoglobulins, lymphocyte surface, isolation and partial characterization, 956
—, suppression of γM, γG, and γA plaque-forming cell responses in cultures of primed spleen cells by class-specific antibody to, 698
—, suppression of primary γM, γG, and γA plaque-forming cell responses in spleen cell cultures by class-specific antibody to, 675
—, synthesis and secretion by chicken lymphoid cells, 1133
Immunolabeling, radio-, technique, sandwich, for lymphocyte surface receptors, quantitative features, 405
Immunologic competence and refractory state in graft-*versus*-host disease in Syrian hamsters, 567
— (IgG) memory in nonresponder mice, selective defect in, 110
— memory cells of bone marrow origin, 1028
Immunological memory and cell-associated immunoglobulin receptors, effect of synthetic polynucleotides on, 45
— memory cells, correlation between binding properties of secreted antibody and cellular receptor for antigen on, 312
— memory cells of thymic origin, 437
— memory to heterologous erythro-

cytes in both T and B cell populations and requirement for T cells in expression of B cell memory, evidence using immunoglobulin allotype and mouse alloantigen θ-markers with congenic mice, 165
Immunologically specific retention of recirculating long-lived lymphocytes in lymph nodes stimulated by xenogeneic erythrocytes, 754
Immunopathogenesis of acute central nervous system disease produced by lymphocytic choriomeningitis virus, 860, 874
Immunosuppression of mice injected with heterologous anti-immunoglobulin heavy chain antisera, 1316
Infection, leukemia virus, murine, of AKR mice, naturally occurring, quantitative studies, 429
—, viral, *in utero*, disease accompanying, role of maternal antibody in tissue injury after transplacental infection with lymphocytic choriomeningitis virus, 827
Inflammation, chronic, in antigen-induced arthritis, pathogenesis, 323
Influenza virus and phytohemagglutinin, receptors for, localization to intramembranous particles, studied by freeze-etching, 1209
Inhibition factor, macrophage migration, production by continuous cell lines, 255
—, selective, of macrophage phagocytic receptors by anti-membrane antibodies, 458
Injection with heterologous anti-immunoglobulin heavy chain antisera, immunosuppression of mice with, 1316
Intracellular fate and ingestion of cholesterol and cholesterol esters, 21
Intramembranous particles, localization of receptors for phytohemagglutinin and influenza virus to, studied by freeze-etching, 1209
Irradiated recipients, sublethally, synergy in graft-*versus*-host reaction quantitated by mortality assay in, 1059

KALLIKREIN, appearance of chemotactic activity for human neutrophils by conversion of human prekallikrein to, 81

LEAD and methyl mercury, effects of acute exposure on cytochrome P-450 and mixed function oxidase system in liver, 1406
Leukemia virus infection, murine, of AKR mice, naturally occurring, quantitative studies, 429
Leukemias and myelomas, murine, viral and cellular surface antigens of, serological analysis by immunoelectron microscopy, 443
Leukocyte cultures, mixed, allogeneic and xenogeneic response in, 1204
Leukocytes, basophilic, in allergic contact dermatitis, 235
—, passenger, role in anomalous survival of neonatal skin grafts, 388
—, polymorphonuclear, proesterase 1, relationship of chemotactic behavior of complement-derived factors, C3a, C5a, and $\overline{C567}$, and bacterial chemotactic factor to ability to activate, 376
Leukosis, EL4, isolation, characterization, and in vitro reactivity of peritoneal lymphoid effector cells from BALB/c mice immune to, 1334
Leukotactic factors elaborated by virus-infected tissues, 1095
Life-span and recirculation of marrow-derived small lymphocytes from thoracic duct, 185
Lipopolysaccharide, *Escherichia coli*, tolerance against, rapid breaking in vivo and in vitro, 850

Listeriosis, requirement of thymus, T, lymphocytes for resistance to, 1104
Liver, effects of acute exposure to lead and methyl mercury on cytochrome P-450 and mixed function oxidase system in, 1406
Lymph nodes stimulated by xenogeneic erythrocytes, immunologically specific retention of recirculating long-lived lymphocytes in, 754
—, plaque-forming cells, ultrastructure, 491
—, thoracic duct, and lymphoid tissue, identification of marrow-derived and thymus-derived small lymphocytes in, 200
Lymphocyte culture, mixed, relationship of response to *HL-A* histocompatibility antigens, effect of allele plus one antigen match, 596
— cytotoxicity in vitro in Moloney sarcoma virus system, synergistic or antagonistic effect of different antibody concentrations on, 997
— migration, thymic, in neonatal mice, magnitude and pattern of, 907
—, sensitized, role in mechanism of fever accompanying delayed hypersensitivity, 1113
— surface immunoglobulins, isolation and partial characterization, 956
— surface receptors, quantitative features of sandwich radioimmunolabeling technique for, 405
—, uropod-bearing, evidence for thymic origin, 1037
Lymphocytes, B, antibody receptor on, detection method and functional significance, 610
—, B, antibody receptor on, immunochemical and electron microscopy characteristics, 627
—, bone–marrow associated, progenitors, quantitative assay for, 1363
—, diploid, synthesis of plasma membrane-associated and secretory immunoglobulin in, 136
Lymphocytes, recirculating long-lived, immunologically specific retention in lymph nodes stimulated by xenogeneic erythrocytes, 754
—, small, marrow-derived, from thoracic duct, life-span and recirculation, 185
—, small, marrow-derived and thymus-derived, identification in lymphoid tissue and thoracic duct lymph, 200
—, T, thymus, requirement for resistance to listeriosis, 1104
—, thymus-derived and marrow-derived, classification by demonstration of antigen-binding characteristics, 1410
—, thymus-derived, radiosensitivity of, in cell-to-cell interaction in immune response, 711
—, thymus-processed, T cells, probable autonomy for killing of allogeneic target cells, 890
— with surface-bound immunoglobulin, ultrastructural study, 267
Lymphocytic choriomeningitis virus, immunopathogenesis of acute central nervous system disease produced by, 860, 874
— choriomeningitis virus, role of maternal antibody in tissue injury after transplacental infection with, 827
Lymphoid cell collaboration, requirement for macrophages in, 1049
— cells, chicken, synthesis and secretion of immunoglobulins by, 1133
— cells, human, nonsecretory, isolation and characterization of cell surface immunoglobulin from, 1392
— cells in delayed hypersensitivity, 985
— cells mediating graft-*versus*-host response, synergy among, 764, 1059
— cells sensitized against allogeneic tumors, specific in vitro deletion and recovery of, 972
— effector cells, peritoneal, from

BALB/c mice immune to EL4 leukosis, isolation, characterization, and in vitro reactivity of, 1334
Lymphoid tissue and thoracic duct lymph, identification of marrow-derived and thymus-derived small lymphocytes in, 200

MACROPHAGE, cholesterol metabolism in, 21
— migration inhibition factor, production by continuous cell lines, 255
— phagocytic receptors, selective inhibition by anti-membrane antibodies, 458
— receptor sites for complement (C3) and for immunoglobulin (IgG), different roles of, 780
Macrophages, requirement for in lymphoid cell collaboration, 1049
Man, genetic polymorphism in glycine-rich β-glycoprotein, 68
—, lymphoid cells, nonsecretory, isolation and characterization of cell surface immunoglobulin from, 1392
—, neutrophils, appearance of chemotactic activity for by conversion of human prekallikrein to kallikrein, 81
—, surface immunoglobulin–bearing tonsil cells, electron microscope study, 533
Mapping of *Ir-1* locus, genetic control of immune response, 1259
Marrow, bone, in vitro proliferation of plasma cells from, role of thymus, 476
— cells, bone, thymocytes from mice immunized against allograft render specifically cytotoxic, 150
— origin, bone, immunologic memory cells of, 1028
Marrow-derived small lymphocytes from thoracic duct, life-span and recirculation, 185
— and thymus-derived lymphocytes, classification by demonstration of antigen-binding characteristics, 1410
Marrow-derived and thymus-derived small lymphocytes, identification in lymphoid tissue and thoracic duct lymph, 200
Maternal antibody, role in tissue injury after transplacental infection with lymphocytic choriomeningitis virus, 827
— antibody to paternal allotype in (SJL × BALB/c)F_1 mice, chronic suppression after perinatal exposure to, 1151
Maternally induced transplantation immunity, tolerance, and runt disease in rats, 808
Membrane, anti-, antibodies, selective inhibition of macrophage phagocytic receptors by, 458
— attack mechanism of complement, molecular analysis, 549
Membrane-associated, plasma, and secretory immunoglobulin, synthesis in diploid lymphocytes, 136
Membranes, erythrocyte, structure studied by freeze-etching, 1209
Memory cells, immunologic, of bone marrow origin, 1028
— cells, immunological, correlation between binding properties of secreted antibody and cellular receptor for antigen on, 312
— cells, immunological, of thymic origin, 437
—, immunologic (IgG), in nonresponder mice, selective defect in, 110
—, immunological, and cell-associated immunoglobulin receptors, effect of synthetic polynucleotides on, 45
—, immunological, to heterologous erythrocytes in both T and B cell populations and requirement for T cells in expression of B cell memory, evidence using immunoglobulin allotype and mouse alloantigen θ-markers with congenic mice, 165
Mercury, methyl, and lead, effects of acute exposure on cytochrome P-

450 and mixed function oxidase system in liver, 1406
Metabolism, cholesterol, in macrophage, 21
— of adenosine 3′,5′-monophosphate in mice exposed to early environmental stress, 220
Microscope, electron, study of surface immunoglobulin–bearing human tonsil cells, 533
Microscopy, electron, and immunochemical characteristics of antibody receptor on B lymphocytes, 627
—, immunoelectron, serological analysis of viral and cellular surface antigens of murine leukemias and myelomas by, 443
Mitogens concanavalin A and phytohemagglutinin, in vitro DNA synthetic response of spleen cells stimulated by, 1279
Molecular analysis of membrane attack mechanism of complement, 549
Molecules, antigen, small mono- and bifunctional, humoral and cellular immune responses to, 1228
Moloney sarcoma virus system, synergistic or antagonistic effect of different antibody concentrations on in vitro lymphocyte cytotoxicity in, 997
Monophosphate, adenosine 3′,5′-, metabolism in mice exposed to early environmental stress, 220
Morphological and functional studies of fetal thymus transplants, 793
Mortality assay in sublethally irradiated recipients, synergy in graft-*versus*-host reaction quantitated by, 1059
Myelomas and leukemias, murine, viral and cellular surface antigens of, serological analysis by immunoelectron microscopy, 443

NEONATAL mice, magnitude and pattern of thymic lymphocyte migration in, 907
— skin grafts, role of passenger leukocytes in anomalous survival of, 388
Nervous system disease, central, acute, produced by lymphocytic choriomeningitis virus, immunopathogenesis, 860, 874
Neutrophils, human, appearance of chemotactic activity for by conversion of human prekallikrein to kallikrein, 81
Nodes, lymph, stimulated by xenogeneic erythrocytes, immunologically specific retention of recirculating long-lived lymphocytes in, 754
Nucleic acid, deoxyribo-, response, synthetic in vitro, of spleen cells stimulated by mitogens concanavalin A and phytohemagglutinin, 1279

ORGAN, contractile, granulation tissue as, structure and function, 719
Oxidase system, mixed function, and cytochrome P-450 in liver, effects of acute exposure to lead and methyl mercury on, 1406

PATERNAL allotype in (SJL × BALB/c)F_1 mice, chronic suppression after perinatal exposure to maternal antibody to, 1151
Pathogenesis of chronic inflammation in antigen-induced arthritis, 323
Perinatal exposure to maternal antibody to paternal allotype in (SJL × BALB/c)F_1 mice, chronic suppression after, 1151
Peritoneal lymphoid effector cells from BALB/c mice immune to EL4 leukosis, isolation, characterization, and in vitro reactivity of, 1334
Phagocytic receptors, macrophage, selective inhibition by anti-membrane antibodies, 458

Phagocytosis of immune complexes by macrophages, 780
Phytohemagglutinin and concanavalin A, mitogens, in vitro DNA synthetic response of spleen cells stimulated by, 1279
— and influenza virus, receptors for, localization to intramembranous particles, studied by freeze-etching, 1209
Picric acid, sensitization with simple chemical compounds, 357
Plaque-forming cell responses, γM, γG, and γA, suppression in cultures of primed spleen cells by class-specific antibody to immunoglobulins, 698
— cell responses, primary γM, γG, and γA, suppression in spleen cell cultures by class-specific antibody to immunoglobulins, 675
— cells of lymph, ultrastructure, 491
Plasma cells, proliferation from bone marrow in vitro, role of thymus, 476
— membrane-associated and secretory immunoglobulin, synthesis in diploid lymphocytes, 136
— prekallikrein, isolation, characterization, and activation mechanism, 1
Polymeric side chains (poly-L-prolyl and poly-DL-alanyl) of synthetic immunogens, contribution of different cell types to genetic control of immune responses as function of chemical nature of, 1009
Polymorphism, genetic, in human glycine-rich β-glycoprotein, 68
Polymorphonuclear leukocytes, proesterase 1, relationship of chemotactic behavior of complement-derived factors, C3a, C5a, and C$\overline{567}$, and bacterial chemotactic factor to ability to activate, 376
Polynucleotides, synthetic, regulation of immune system by, 45
Polypeptide chains, immunoglobulin, idiotypic specificity as potential genetic marker for variable regions of, 579
Prealbumin activator of prekallikrein, 81
Prekallikrein, human, appearance of chemotactic activity for human neutrophils by conversion to kallikrein, 81
—, plasma, isolation, characterization, and activation mechanism, 1
Proesterase 1 of polymorphonuclear leukocytes, relationship of chemotactic behavior of complement-derived factors, C3a, C5a, and C$\overline{567}$, and bacterial chemotactic factor to ability to activate, 376
Protein conjugates, hapten, induction of immunity and tolerance in vitro by, 735

RADIOIMMUNOLABELING technique, sandwich, for lymphocyte surface receptors, quantitative features, 405
Radiosensitivity of thymus-derived lymphocytes in cell-to-cell interaction in immune response, 711
Receptor, antibody, on B lymphocytes, detection method and functional significance, 610
—, antibody, on B lymphocytes, immunochemical and electron microscopy characteristics, 627
—, cellular, for antigen on immunological memory cells, and secreted antibody, correlation between binding properties, 312
— sites, macrophage, for complement (C3) and for immunoglobulin (IgG), different roles of, 780
Receptors for phytohemagglutinin and influenza virus, localization to intramembranous particles, studied by freeze-etching, 1209
—, immunoglobulin, cell-associated, and immunological memory, effect of synthetic polynucleotides on, 45
—, lymphocyte surface, quantitative

features of sandwich radioimmunolabeling technique for, 405
—, macrophage phagocytic, selective inhibition by anti-membrane antibodies, 458
— on immunocompetent cells, cellular correlates of maturation of immune response, 660
— on immunocompetent cells, direct measurement of avidity and cooperative binding of multivalent ligands, 643
Resistance to listeriosis, requirement of thymus, T, lymphocytes for, 1104
Ribonucleic acid, deoxy-, response, synthetic in vitro, of spleen cells stimulated by mitogens concanavalin A and phytohemagglutinin, 1279
Runt disease, maternally induced transplantation immunity, and tolerance in rats, 808

SARCOMA virus system, Moloney, synergistic or antagonistic effect of different antibody concentrations on in vitro lymphocyte cytotoxicity in, 997
— virus–transformed BALB/3T3 nonproducer cells, murine, antigenic properties of, 503
Scleroderma skin fibroblasts in cell culture, connective tissue synthesis by, 1351
Secretion and synthesis of immunoglobulins by chicken lymphoid cells, 1133
Secretory, non-, human lymphoid cells, isolation and characterization of cell surface immunoglobulin from, 1392
— and plasma-membrane associated immunoglobulin, synthesis in diploid lymphocytes, 136
Sensitivity, radio-, of thymus-derived lymphocytes in cell-to-cell interaction in immune response, 711
Sensitization with simple chemical compounds, picric acid, 357
Sensitized lymphocyte, role in mechanism of fever accompanying delayed hypersensitivity, 1113
Sera, anti-I and anti-i, classification into groups based on reactivity patterns with various antigens related to blood group A, B, H, Le^a, Le^b and precursor substances, 1247
Serological analysis by immunoelectron microscopy of viral and cellular surface antigens of murine leukemias and myelomas, 443
Serologically demonstrable alloantigens of epidermal cells, 938
Skin fibroblasts, scleroderma, in cell culture, connective tissue synthesis by, 1351
— grafts, allogeneic, cell-mediated cytotoxicity during rejection and enhancement of, 1301
— grafts, neonatal, role of passenger leukocytes in anomalous survival of, 388
Specific, class-, antibody to immunoglobulins, suppression of γM, γG, and γA plaque-forming cell responses in cultures of primed spleen cells by, 698
—, class-, antibody to immunoglobulins, suppression of primary γM, γG, and γA plaque-forming cell responses in spleen cell cultures by, 675
— heterologous enhancement of immune responses, 437
— immune response genes, GA and GT, influence on specificity of cellular and humoral immune responses to terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, 98
— in vitro cytotoxicity, cells mediating, 890
— in vitro deletion and recovery of lymphoid cells sensitized against allogeneic tumors, 972
— retention, immunologically, of recirculating long-lived lymphocytes in lymph nodes stimulated by xenogeneic erythrocytes, 754
Specificities, idiotypic, suppression by administration of antiidiotypic antibody, 1293

Specificity, idiotypic, as potential genetic marker for variable regions of immunoglobulin polypeptide chains, 579
— of cellular and humoral immune responses to terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, influence of GA and GT immune response genes on, 98
Spleen cell cultures, primed, suppression of γM, γG, and γA plaque-forming cell responses in by class-specific antibody to immunoglobulins, 698
— cell cultures, suppression of primary γM, γG, and γA plaque-forming cell responses in by class-specific antibody to immunoglobulins, 675
— cells stimulated by mitogens concanavalin A and phytohemagglutinin, in vitro DNA synthetic response of, 1279
— cells, transfer of suppressing factor with, active suppression of immunoglobulin allotype synthesis, 1163
Stimulation of spleen cells by mitogens concanavalin A and phytohemagglutinin, in vitro DNA synthetic response, 1279
Synergistic or antagonistic effect of different antibody concentrations on in vitro lymphocyte cytotoxicity in Moloney sarcoma virus system, 997
Synergy among lymphoid cells mediating graft-*versus*-host response, 764, 1059

TARGET cells, allogeneic, probable autonomy of thymus-processed lymphocytes, T cells, for killing of, 890
Terpolymer of L-glutamic acid, L-alanine, and L-tyrosine, influence of GA and GT immune response genes on specificity of cellular and humoral immune responses to, 98
Thoracic duct, life-span and recirculation of marrow-derived small lymphocytes from, 185
— duct lymph and lymphoid tissue, identification of marrow-derived and thymus-derived small lymphocytes in, 200
Thymectomy, effect on primary and secondary antibody response to poly-L(Tyr, Glu)-poly-D, L-Ala–poly-L-Lys, 126
Thymic lymphocyte migration in neonatal mice, magnitude and pattern of, 907
— origin, immunological memory cells of, 437
— origin of uropod-bearing lymphocyte, evidence for, 1037
Thymocytes from mice immunized against allograft render bone marrow cells specifically cytotoxic, 150
Thymus function, duration, 339
—, role in in vitro plasma cell proliferation from bone marrow, 476
—, T, lymphocytes, requirement for resistance to listeriosis, 1104
— transplants, fetal, morphological and functional studies, 793
Thymus-derived cell function, genetic control of, 1279
— cells, two types, evidence for interaction between in graft-*versus*-host response, 764
— lymphocytes, radiosensitivity of, in cell-to-cell interaction in immune response, 711
— and marrow-derived lymphocytes, classification by demonstration of antigen-binding characteristics, 1410
— and marrow-derived small lymphocytes, identification in lymphoid tissue and thoracic duct lymph, 200
Thymus-processed lymphocytes, T cells, probable autonomy for killing of allogeneic target cells, 890
Tissue, connective, synthesis by scleroderma skin fibroblasts in cell culture, 1351

Tissue, granulation, as contractile organ, structure and function, 719
— injury after transplacental infection with lymphocytic choriomeningitis virus, role of maternal antibody in, 827
—, lymphoid, and thoracic duct lymph, identification of marrow-derived and thymus-derived small lymphocytes in, 200
Tissues, collagenous, preferential localization of antigen-antibody complexes to in antigen-induced arthritis, 323
—, virus-infected, leukotactic factors elaborated by, 1095
Tolerance against *Escherichia coli* lipopolysaccharide, rapid breaking in vivo and in vitro, 850
—, antibody, high and low zone, to flagellin, enhanced cell-mediated immunity during, 298
— and immunity, induction in vitro by hapten protein conjugates, 735
—, maternally induced transplantation immunity, and runt disease in rats, 808
— of living cells, immune status of mice, 1185
Tonsil cells, human, surface immunoglobulin-bearing, electron microscope study, 533
Transplacental infection with lymphocytic choriomeningitis virus, role of maternal antibody in tissue injury after, 827
Transplantation immunity, cellular immunoabsorbents in, 972
— immunity, maternally induced, tolerance, and runt disease in rats, 808
Transplants, thymus, fetal, morphological and functional studies, 793
Trophoblast, autoantigenic status, further evidence concerning, 1177
Tumor allografts, ascites, rejection, 1334
— cells, antigen suppression in, two kinds revealed by cell fusion, 839
Tumors, allogeneic, specific in vitro deletion and recovery of lymphoid cells sensitized against, 972
L-Tyrosine, L-glutamic acid, and L-alanine, terpolymer of, influence of GA and GT immune response genes on specificity of cellular and humoral immune responses to, 98

UROPOD-BEARING lymphocyte, evidence for thymic origin, 1037

VIRAL and cellular surface antigens of murine leukemias and myelomas, serological analysis by immunoelectron microscopy, 443
— infection, *in utero*, disease accompanying, role of maternal antibody in tissue injury after transplacental infection with lymphocytic choriomeningitis virus, 827
Virus, choriomeningitis, lymphocytic, role of maternal antibody in tissue injury after transplacental infection with, 827
— infection, leukemia, murine, of AKR mice, naturally occurring, quantitative studies, 429
—, influenza, and phytohemagglutinin, receptors for, localization to intramembranous particles, studied by freeze-etching, 1209
—, lymphocytic choriomeningitis, immunopathogenesis of acute central nervous system disease produced by, 860, 874
— system, Moloney sarcoma, synergistic or antagonistic effect of different antibody concentrations on in vitro lymphocyte cytotoxicity in, 997
Virus-infected tissues, leukotactic factors elaborated by, 1095

Virus-transformed BALB/3T3 nonproducer cells, murine sarcoma, antigenic properties, 503

XENOGENEIC and allogeneic response in mixed leukocyte cultures, 1204
— erythrocytes, immunologically specific retention of recirculating long-lived lymphocytes in lymph nodes stimulated by, 754
X-Irradiation, influence on passive transfer and on in vitro production of soluble mediators, lymphoid cells in delayed hypersensitivity, 985